



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Another male (No. 63,621, Smiths. Inst.), moulting, but in nearly fresh plumage, differs from the above in being much less heavily barred with black on the back and scapulars, and of a still deeper shade of tawny ochraceous below.

Description of female type.—Crown bluish slate, streaked with chestnut rufous on the vertex. Above chestnut rufous, sparingly barred with plumbeous black, the bars becoming obsolete on the rump, leaving the upper tail coverts and hinder part of rump plain; tail incompletely barred with black, the outer feathers only spotted on their inner webs. Underparts pale ochraceous buff, deepening to vinaceous cinnamon on the breast; longitudinal markings below light cinnamon, nearly obsolete.

Measurements.—Wing, 200.0; tail, 152.0; chord of culmen, 13.0; width of bill at base, 10.5; tarsus, 39.0; middle toe, 24.5 mm.

DESCRIPTION OF AN APPARENTLY NEW *CENTURUS* FROM GREAT BAHAMA ISLAND, BAHAMAS.

BY CHARLES B. CORY.

A COLLECTION of some eight hundred birds lately received from Great Bahama Island contained eight specimens of an apparently undescribed *Centurus* which I propose to call

Centurus bahamensis, SP. NOV.

Type No. 19,578, Coll. C. B. Cory, Boston, Great Bahama Island, December 26, 1891. — Adult male similar to *C. blakei*, in having the entire underparts strongly tinged with olive green; the back is banded with black and yellowish green, not black and white as in *blakei*. The feathers on the flanks show a slightly reddish tinge; the forehead is dusky white; and the red at the base of the bill is somewhat darker than in *C. blakei*. Length, 9.50 inches; wing, 5.25; tail, 4.00; bill, 1.20; tarsus, .85.

The female has the forehead darker than the female of *C. blakei*, and has the under parts tinged with olive green as in the male.